



---

---

## "EFFECT OF OXYTETRACYCLIN ON BEHAVIOR OF FISH CATLA CATLA"

**Dr. Dinesh Prasad Patel, Dr. Vinod Kumar Mishra and Pramod Kumar Prajapati**

**Assistant Professor , Department of Zoology ,  
Govt. S.K. College Mauganj, Distt. – Rewa (M.P.).**

### ABSTRACT

*The present study was designed to assess the toxic effect of antibiotic oxytetracyclin in fish Catla catla using behavior changes as a biomarker. The acute exposure was given up to 96 hours of treatment. Sub-lethal concentration used in the experiment was 80 mg/l of water. Change in behaviour pattern was observed after 24, 48, 72 and 96 hours of exposure. Different behavioural alteration like, high pigmentation, high mucus secretions, loss of equilibrium, hyperactivity were seen throughout the exposure. The behavioural changes become more severe with increase in time duration.*



**KEYWORDS:** Behaviour, toxicity, antibiotics and oxytetracyclin.

### INTRODUCTION

Oxytetracyclin is widely used broad spectrum antibiotic belonging to class tetracycline (Li W et al., 2012). In a world market of pharmaceuticals India has emerged as one of the top five pharmaceuticals market (Chander V et al., 2016). Production of pharmaceuticals is rising at 14% per year. This may be due to excise duty free zone for pharmaceutical manufacturing done by the Central and State Governments of India leading to large scale production of pharmaceutical. Moreover deliberate discharge and accidental release of non biodegradable by-products either untreated or partially treated waste to environment further deepen the problem of pharmaceutical pollution in aquatic ecosystems and has the potential to disrupt the structure and functioning of natural ecosystems.

Oxytetracyclin has high hydrophilic characters and low volatilization and significantly persistent in water (Daghrir R et al., 2013). Bioassay based on behaviour is faster, more sensitive and ecologically more relevant as assessing growth and reproduction need longer bioassay. Behavioural bioassay is more promising alternatives than lethality evaluating bioassay which are currently used for the risk assessment of toxicant. Behavioural changes provide early warning signals about the health of exposed population which other standard tests do not take in to consideration (Sharma M et al., 2015). Fish are very sensitive to even minute quantities of pollutants so can be considered sentinels of slight perturbation in the environment. Furthermore, fish is a good indicator of toxicity studies due to its position in food web ability to bio-accumulate toxicants sensitivity to low concentration of pollutant and nutritive value to human (Sharma M et al., 2019). Fish *Catla catla* was used for the experiment. So the present study was designed to study the toxic effect of oxytetracyclin on behaviour of fish.

**MATERIALS AND METHODS:**

The specimens of experimental fish *Catla catla* were collected from Govindgarh lakes of district Rewa. The chosen fish were of total length  $23.5 \pm 17$  cm and total weight of  $160 \pm 60$  g. The antibiotic oxytetracyclin immersion was taken from the local medical store. Exposure was semi-static and every 24 hours the drug was replenished to complete the initial concentration. The experimental glass aquaria were of 100 liters capacity each. Control animal were submitted to the same water change schedule without the addition of Oxytetracyclin. Exposure was given for 96 hours. Sub-lethal concentration of Oxytetracyclin was decided as 80 mg/l of water for the exposure according to the LC50 value for *Catla catla* of Ambili (Ambili TR et al., 2013).

Behavioural alterations were observed after 24, 48, 72 and 96 hours of exposure from both treated as well as control groups. Behavioural inconsistencies were keenly observed in both the groups (Control and treated) for hyperactivity, movement, equilibrium, opercula movement, pigmentation, respiratory rate, air gulping, jumping, mucus secretion, aggressiveness, motionless, vertical movement, surfacing.

**RESULTS:**

Control or untreated fish exhibited normal behavior like a normal rate of swimming, mild opercular movement, responding to low stimulus during the experiment. On the other hand fish exposed to Oxytetracyclin, significant changes in behaviour were observed. As the fish were placed in the exposure medium they immediately move to the corner of the tank. But after a few hours they become mild hyperactive and move separately in the tank. After 24 hours of the exposure, mild increase in pigmentation was observed and a layer of mucus was also seen on the body surface. With increase in duration of exposure surfacing behaviour was observed along with hyperventilation, loss of equilibrium, high pigmentation, thick layer of mucus, very less movement and lost response to external stimuli (Table-1). At the highest hour of exposure pigmentation, loss of equilibrium, mucus secretion and surface gulping was more pronounced.

**Table 1: Different behavioural changes observed in fish *Catla catla* after exposure with 80 mg/liters of Oxytetracyclin.**

S.N.	Parameters	Control	24 hrs	48 hrs	72 hrs	96 hrs
1	Hyperactivity	-	+	++	++	++
2	Operculum movement	+	--	++	++	++
3	Rate of swimming	+	--	+	++	++
4	Corner behaviour	-	+++	-	-	-
5	Pigmentation	-	+	+++	+++	+++
6	Mucus secretion	-	+	++	++	++
7	Loss of balance	-	-	-	+	++
8	Surface gulping	-	-	+	+++	+++

Where: (-) None, (+) mild, (++) Moderate, (+++) Strong

**DISCUSSION:**

Behaviour analysis may be considered as a robust and sensitive tool for detecting acute exposure of chemical to fish (Gaaied S et al., 2019). Behavioural alterations may be a protective mechanism of fish to avoid unsuitable conditions for some time or may be toxic damage to the nervous system. In the present study the locomotor response of fish was sensitive to Oxytetracyclin as observed by alteration in the swimming pattern of fish. Alteration of locomotion behaviour is very often linked with neurological impairment. Tetracyclins have been associated with cranial nerve toxicity and neuromuscular blockage (Thomas RJ et al., 1991). Respiratory distress observed may be due to increase in metabolic demand of fish. Decline in swimming activity with increase in exposure time may be a result of the increase demand of glycogen to meet the stressful condition. Aquatic surface respiration

was observed at later hour of exposure due to enhanced oxygen demand due to hypoxic condition. Excessive mucus secretion found in fish is to make a protective layer around the body to reduce the body contact toxic environment and get relief from irritation caused by pollutant (Sharma M 2019). A high pigmentation was observed, which suggests that Oxytetracyclin dysfunction the pituitary gland and influence the distribution of chromatophores (Harit G & Srivastava N 2018). Previous studies have revealed that a number of chemicals can provoke abnormal behavior in fish (Sharma M 2019 & Altenhofen S et al., 2017). Several studies have reported similar results in fish with response to pharmaceuticals (Almeida AR et al., 2019).

## CONCLUSION:

The results of the present show that oxytetracyclin has a strong influence on behavior of *Catla catla* and contribute to sound assessment of environmental hazards posed by this antibiotic.

## REFERENCES:

- Almeida AR, Tacao M, Machado AL, Golovko O, Zlabek V, Domingues I, et al. Long-term effects of oxytetracycline exposure in zebrafish: A multi-level perspective. *Chemosphere* 2019;222:333-344.
- Altenhofen S, Wiprich MT, Nery LR, Leite CE, Vianna MR, Bonan CD, et al. Manganese(II) chloride alters behavioral and neurochemical parameters in larvae and adult zebrafish. *Aquatic Toxicology* 2017;182:172-183. <https://doi.org/10.1016/j.aquatox.2016.11.013>
- Ambili TR, Saravanan M, Ramesh M, Abhijith DB, Poopal RK. Toxicological effects of the antibiotic oxytetracycline to an Indian major carp *Labeo rohita*. *Archive of Environmental Contamination and Toxicology* 2013;64(3):494-503. doi: 10.1007/s00244-012-9836-6. Epub 2012 Nov 29.
- Chander V, Sharma B, Negi V, Aswal RS, Singh P, Singh R, et al. Pharmaceutical Compounds in Drinking Water. *Journal of Xenobiotics* 2016;6(1):5774. <http://doi:10.4081/xeno.2016.5774>
- Daghrir R, Drogui P. Tetracycline antibiotics in the environment: a review. *Environmental Chemistry Letters* 2013;11(3):209-227.
- Gaaied S, Oliveira M, Domingues I, Banni M. 2,4- Dichlorophenoxyacetic acid herbicide effects on zebrafish larvae: development, neurotransmission and behavior as sensitive endpoints. *Environmental Science and Pollution Research* 2019; <https://doi.org/10.1007/s11356-019-04488-5>
- Harit G, Srivastava N. Behavioural alterations in *Channa punctatus* after exposure to endosulfan followed by subsequent recovery. *International Journal of Fisheries and Aquatic Studies* 2018;6(2):51-55
- Li W, Shi Y, Gao L, Liu J, Cai Y. Occurrence of antibiotics in water, sediments, aquatic plants, and animals from Baiyangdian Lake in North China. *Chemosphere* 2012;89(11):1307-1315
- Sharma M, Chadha P, Borah MK. Fish behaviour and immune response as a potential indicator of stress caused by 4-nonylphenol. *American Journal of Biosciences* 2015;3(6):278-283.
- Sharma M, Chadha P. Immunological impairment as a marker of 4-nonylphenol induced stress. *Indian Journal of Animal Sciences* 2019;89(6):695-698, June 2019/Article
- Sharma M. Behavioural responses in effect to chemical stress in fish: A review. *International Journal of Fisheries Aquatic Studies* 2019;7(1):01-05.
- Thomas RJ. Neurotoxicity of antibacterial therapy. *Southern Medical Journal* 1991;87:869-74.



**Dr. Dinesh Prasad Patel**  
**Assistant Professor , Department of Zoology ,**  
**Govt. S.K.N. College Mauganj, Distt. – Rewa (M.P.)**