

Vol II Issue X July 2013

ISSN No : 2249-894X

*Monthly Multidisciplinary
Research Journal*

*Review Of
Research Journal*

Chief Editors

Ashok Yakkaldevi
A R Burla College, India

Flávio de São Pedro Filho
Federal University of Rondonia, Brazil

Ecaterina Patrascu
Spiru Haret University, Bucharest

Kamani Perera
Regional Centre For Strategic Studies,
Sri Lanka

Welcome to Review Of Research

RNI MAHMUL/2011/38595

ISSN No.2249-894X

Review Of Research Journal is a multidisciplinary research journal, published monthly in English, Hindi & Marathi Language. All research papers submitted to the journal will be double - blind peer reviewed referred by members of the editorial Board readers will include investigator in universities, research institutes government and industry with research interest in the general subjects.

Advisory Board

Flávio de São Pedro Filho Federal University of Rondonia, Brazil	Horia Patrascu Spiru Haret University, Bucharest, Romania	Mabel Miao Center for China and Globalization, China
Kamani Perera Regional Centre For Strategic Studies, Sri Lanka	Delia Serbescu Spiru Haret University, Bucharest, Romania	Ruth Wolf University Walla, Israel
Ecaterina Patrascu Spiru Haret University, Bucharest	Xiaohua Yang University of San Francisco, San Francisco	Jie Hao University of Sydney, Australia
Fabricio Moraes de Almeida Federal University of Rondonia, Brazil	Karina Xavier Massachusetts Institute of Technology (MIT), USA	Pei-Shan Kao Andrea University of Essex, United Kingdom
Catalina Neculai University of Coventry, UK	May Hongmei Gao Kennesaw State University, USA	Osmar Siena Brazil
Anna Maria Constantinovici AL. I. Cuza University, Romania	Marc Fetscherin Rollins College, USA	Loredana Bosca Spiru Haret University, Romania
Romona Mihaila Spiru Haret University, Romania	Liu Chen Beijing Foreign Studies University, China	Ilie Pintea Spiru Haret University, Romania
Mahdi Moharrampour Islamic Azad University buinzahra Branch, Qazvin, Iran	Nimita Khanna Director, Isara Institute of Management, New Delhi	Govind P. Shinde Bharati Vidyapeeth School of Distance Education Center, Navi Mumbai
Titus Pop PhD, Partium Christian University, Oradea, Romania	Salve R. N. Department of Sociology, Shivaji University, Kolhapur	Sonal Singh Vikram University, Ujjain
J. K. VIJAYAKUMAR King Abdullah University of Science & Technology, Saudi Arabia.	P. Malyadri Government Degree College, Tandur, A.P.	Jayashree Patil-Dake MBA Department of Badruka College Commerce and Arts Post Graduate Centre (BCCAPGC), Kachiguda, Hyderabad
George - Calin SERITAN Postdoctoral Researcher Faculty of Philosophy and Socio-Political Sciences Al. I. Cuza University, Iasi	S. D. Sindkhedkar PSGVP Mandal's Arts, Science and Commerce College, Shahada [M.S.]	Maj. Dr. S. Bakhtiar Choudhary Director, Hyderabad AP India.
REZA KAFIPOUR Shiraz University of Medical Sciences Shiraz, Iran	Anurag Misra DBS College, Kanpur	AR. SARAVANAKUMARALAGAPPA UNIVERSITY, KARAIKUDI, TN
Rajendra Shendge Director, B.C.U.D. Solapur University, Solapur	C. D. Balaji Panimalar Engineering College, Chennai	V.MAHALAKSHMI Dean, Panimalar Engineering College
	Bhavana vivek patole PhD, Elphinstone college mumbai-32	S.KANNAN Ph.D , Annamalai University
	Awadhesh Kumar Shirotriya Secretary, Play India Play (Trust), Meerut (U.P.)	Kanwar Dinesh Singh Dept.English, Government Postgraduate College , solan

More.....

Address:-Ashok Yakkaldevi 258/34, Raviwar Peth, Solapur - 413 005 Maharashtra, India
Cell : 9595 359 435, Ph No: 02172372010 Email: ayisrj@yahoo.in Website: www.isrj.net



STUDIES ON COLD TREATMENT EFFECT ON GERMINATION OF BRASSICA JUNCEA SEEDS

S.S.PHULARI

Principal , MSPM's D. G. TATKARE COLLEGE
(Arts, Science, Commerce) Mangaon Raigad, Maharashtra, India.

Abstract:

The Brassica juncea seeds soaked in water for 24 hours were treated with cold temperature 20, 40, 60 and 80 minutes before sowing. The observation were made daily upto 10 days of sowing with adequate watering. There was no significant variation in control and cold treated seeds with respect to days for germination, germination % vigor %, plumule formation, average shoot length, average number of roots, root length and fresh weight of seedlings. The days for germination in control and cold treated seeds was about the same. The germination % and vigor % not affected by cold treatment. Plumule formation average shoot length in cold treated seeds was neither increased or decreased over control. The root number, root length and fresh weight of seedling neither stimulating nor suppressing due to cold treatment as compared to control.

KEY WORDS:

Cold Treatment , Germination ,brassica Juncea Seeds , Materials And Methods .

.INTRODUCTION

Brassica juncea , also known as mustard greens, Indian mustard and leaf mustard. Leaves, seeds and stems are used in in Indian cuisine (Chandrasekaran, 2013). Indian Mustard is a folk remedy for arthritis, footache , lumbago and rheumatism (Duke and Wain, 1981). Chinese eat the leaves in soups for bladder, inflammation or haemorrhage. Mustard oil is used for skin eruptions and ulcers (Perry, 1980). Leaves applied to the forehead are said to relieve headache (Burkill, 1966). Many species of plants are capable of adapting to low temperature and freezing conditions. Generally, this adaptations requires a prior period of acclimation at low, non freezing temperatures, during which time a number of morphological, physiological and molecular changes occur (Singh and Laroche, 1988, Guy, 1990; Alberdi and (Cocuera, 1991). In present investigation, the Brassica juncea seeds are treated with cold temperature (0 c) for 20 minutes, 40 minutes, 60 minutes and 80 minutes. The Brassica juncea pregerminated seeds treated with cold temperature and sown to study the effect of cold temperature on the germination. The germination is vital process. The cold treatment may affects the germination percentage, plumule formation, short length, root number, roof length and fresh weight of seedling. The physiological changes may reflect in the morphology of germination.

MATERIALS AND METHODS :-

Seeds of Brassica juncea were kept in petridish containing moist blotting papers. The Petridish containing seeds kept under dark condition for 24 hrs. It triggers the process of germination. The germinations ready seeds were subjected to cold temperature (0 C) for 20 minute, 40 minute, 60 minute and 80 minute. For each treatment, 10 seeds were subjected. The treated and control seeds then sown in

Title: STUDIES ON COLD TREATMENT EFFECT ON GERMINATION OF BRASSICA JUNCEA SEEDS
Source: Review of Research [2249-894X] S.S.PHULARI yr:2013 vol:2 iss:10

pots containing garden mixture. The process is done in two replica.

The sown seeds were watered adequately. The observations were made considering the parameters of germination such as germination percentage, days required for germination, days required to emergence of plumage, length of seedling, number of roots per seedling, average length of root per seedling, and fresh weight of seedling. The experiment was set in two replicas. In each replica for each treatment ten pregerminated Brassica juncea seeds were selected. The observation were made for ten days from date of sowing the seeds. Plants were uprooted on 10th day of sowing after treatment for number of roots length of roots and fresh weight of seedling. The average is work out by applying arithmetic mean biostatistical application.

Germination was deemed to have occurred when the radical was visibly extended beyond the surface of the seed, usually protruding through a fracture in the seed coat. Number of seeds germinated each day was counted and % of germination was calculated. Germination vigor was determined using the method described by (Zabator(1962), modified to include the number of days of incubation at each temperature and each treatment. modified to include the number of days of incubation at each temperature and each treatment.

The vigor value was calculated as follows :- $V = (a/1 + b/2 + c/3 + \dots + x/n)/s \times 100$ Where a,b,c,d and x represent the number of seeds which had germinated after the first, second, third, fourth and nth day of incubation n respectively, and "s" represents the total number of seeds in the sample.

RESULTS AND DISCUSSION :-

(I) Effect of cold treatment on germination of Brassica juncea seeds. Effect of cold temperature treatment on germination of Brassica juncea seeds.

Days after sowing	Cold Treatment (germination out of 10)				
	Control	20 min.	40 min	60 min	80 min
1 st	-	-	-	-	-
2 nd	-	-	-	-	-
3 rd	2	2	2	4	3
4 th	3	2	4	4	4
5 th	4	5	4	5	4
6 th	6	5	6	5	5
7 th	8	7	7	8	7
8 th	8	8	7	8	9
9 th	9	9	9	10	10
10 th	10	10	10	10	10

Germination temperatures vary greatly among plants. A plants natural environment and more specifically the environmental conditions at the time of natural seed disposal, will reveal information regarding the optimal temperature range for germination. Alternating temperature regimes are often more favorable for germination than constant temperatures, Some species will only germinates at alternating temperature. (Baskin and Baskin 2001).

Table I shows the number of seeds germinated out of ten seeds on the days after sowing. There is no germination of seeds for first and second day of sowing. The control and cold treated seeds of Brassica juncea germinated since the 3rd day of sowing. There is no any significant difference between number of seeds germinated in control and cold treated seeds. On 10th day at sowing, all seeds germinated of control as well as 20 min, 40 min, 60 min and 80 min. cold temperature treated seeds of Brassica, Bellairs and Bell (1990) reported that low temperatures tended not to affect germination. According to Khan and Ungar (1999) at higher temperature differences between light and dark germinated seeds were not significant.

(I) Effect of cold treatment on germination % of Brassica juncea seeds :- Table II Effect of cold treatment on Brassica juncea seeds germination

Days after sowing	Cold Treatment germination %				
	Control	20 min.	40 min	60 min	80 min
3 rd	20 %	20 %	20 %	40 %	30 %
4 th	30 %	20 %	40 %	40 %	40 %
5 th	40 %	50 %	40 %	50 %	40 %
6 th	60 %	50 %	60 %	50 %	50 %
7 th	80 %	70 %	70 %	80 %	70 %
8 th	80 %	80 %	70 %	80 %	90 %
9 th	90 %	90 %	90 %	100 %	100 %
10 th	100 %	100 %	100 %	100 %	100 %

Seed germination depends on external factors include temperature, water, oxygen. Various plants require different variables for successful seed germination. The temperature may induce or inhibit the seed germination. To study the effect of cold treatment on seed germination may prove a significant factor.

Table II indicates the effects of cold treatment germination % of Brassica juncea seeds. It seems to be about 20% to 40% seed germination in control and cold-treated seeds on the 5th day of sowing. The 100% germination of seeds is seen on the 10th day in control and cold-treated seeds of Brassica juncea seeds. There is no significant variation in the germination percentage of the control and 20 min, 40 min, 60 min and 80 min cold-treated seeds. A 2009 study of Carroll Sarah, showed that temperature has a significant effect on both the time to germinate and the proportion of mustard seeds. However, the present study reveals that there is no significant variation in germination due to cold temperature treatment. It is concluded from the above results and discussion that the germination percentage is not affected by cold treatment in Brassica juncea seeds.

(I) Effect of cold treatment on vigor value of Brassica juncea seeds :-

Table III – Effect of cold treatment on vigor value of Brassica juncea seeds :-

Control	20 min.	40 min	60 min	80 min
0.0076	0.0069	0.0073	0.0074	0.0071

Table III shows the effect of cold treatment on the vigor value of Brassica juncea seeds. From the Table III it is clear that the vigor value of control seeds is 0.0076. The vigor value of cold-treated seeds of Brassica juncea is in the range of 0.0069 to 0.0074. Thus, there is no significant variation in the vigor value of control seeds as well as cold-treated seeds of Brassica juncea. High temperature affects the yield and quality of Brassica juncea. (Anan et al. 2010). The plants of Brassica juncea exposed to temperature stress exhibited a significant decline in growth (Hayat et al. 2000).

(I) Effect of cold treatment on plumule formation in Brassica juncea seeds :-

Table IV – Effect of cold treatment on plumule formation in Brassica juncea seeds

Days from sowing	Cold Treatment (Days required for plumule formation)				
	Control	20 min.	40 min	60 min	80 min
4 th day	-	-	-	-	-
5 th day					

Table IV shows the effect of cold treatment on plumule formation in Brassica juncea seeds germination. It is observed that in control as well as cold-treated seeds of Brassica juncea show plumule formation on the fifth day of sowing. There is no variation in plumule formation during germination in Brassica seeds due to cold temperature.

(V) Effect of cold treatment on average length of shoot in Brassica juncea seeds:-**Table V – Effect of cold treatment on av. Length of shoot in Brassica juncea seeds.**

Days after sowing	Control	20 min.	40 min	60 min	80 min
10 th day	1.7 cm	1.6 cm	1.6 cm	1.5 cm	1.6 cm

Table V depicts the effect of cold treatment on average length of shoot in Brassica Juncea seed germination. On the tenth day, the average length of seedling in control and cold treated seeds is almost similar. There is 1.6 cm average seedling length in control seedling of Brassica juncea. However the length of seedling is in between 1.5 cm to 1.6 cm. in the cold treated seeds at 20 min, 40 min, 60 min and 80 minutes. There is no remarkable change in the seedling length due to cold treatment over the control.

(VI) Effect of cold treatment during germination on number of roots and root length in Brassica juncea:-**Table VI Effect of cold treatment during germination on number of roots, in Brassica Juncea seeds and root length**

Days after Sowing	Control		Cold of 20 min.		Cold of 40 min		Cold of 60 min		Cold of 80 min	
	Number Of roots Per seeding	Root Length cm	Number Of roots Per seeding	Root Length cm	Number Of roots Per seeding	Root Length cm	Number Of roots Per seeding	Root Length cm	Number Of roots Per seeding	Root Length cm
10 th day	3.1	1.1	2.8	1.4	3.2	1.0	3.3	1.2	3.6	1.3

The table VI indicates the effect of cold treatment during germination on number of roots and root length in Brassica juncea seeds. It is clear from the table that average number of roots per seedling is 3.1 in control seedling. The average number of roots per seedling is in range of 2.8, 3.2, 3.3 and 3.6 in Brassica juncea seeds treated with 20 min, 40 min, 60 min and 80 min cold treatment respectively. It seems to be no remarkable change in average root length in seedling due to cold treatment for variable time period. The table VI also shows the average root length per seedling in control as well as cold treated seeds of Brassica juncea. The root length is 1.1 cm in control seedling while it is 1.4 cm i.e. maximum root length in 20 minutes cold treated seedlings. The 40 min, 60 min & 80 minutes treated seedlings shows the root length as 1.0 cm, 1.2 cm and 1.3 cm respectively. Thus, it is observed no consistent variation in the root length in control and cold treated seedlings.

(VI) Effect of cold treatment on fresh weight during germination of seedlings in Brassica Juncea :-**Table VII- Effect of cold treatment on fresh weight during germination of seedlings in Brassica juncea.**

Days after sowing	Cold Treatment				
	Control (gm.)	20 min. (gm.)	40 min (gm.)	60 min (gm.)	80 min (gm.)
10 th day	0.56	0.53	0.52	0.54	0.51

Table shows the effect of cold treatment on fresh weight of seedling during germination of Brassica juncea seeds on 10th day of sowing. The range of fresh weight of 10 seedlings in control and cold treated seeds is 0.51 gm to 0.56 gm. The variation in fresh weight due to cold treatment over the control is

no remarkable. It is non significant change.

BIBLIOGRAPHY :-

- 1) Alberdi, K; Cocuera, L.J. 1991. Cold acclimation in plants. *Phytochemistry*, 30:3177- 3184
- 2) Anand Anjali; Nagarajan, S; Kishore, N; and Verma, A.P.S. 2010. *The Indian J of Agri Sc.* 80 (12).
- 3) Baskin, C. and Baskin, J. 2001. *Seeds, Ecology, Biogeography and Evaluation of Dormancy and germination*, London, Academic Press.
- 4) Bellairs, S.M. and Bell, D.T. (1990) Temperature effects on the seed-germination of 10 Kwongan species from Eneabba, Western-Australia, *American Journal of Botany* 38 (5) 451-458.
- 5) Burkill, J.H. 1966. *A dictionary of economic products of the Malay Peninsula*. Art Printing works, Kuala Lumpur, 2 vols.
- 6) Carroll, S. 2004. The effect of temperature on the germination of mustard seeds. *Dublin Magazine. The Sunday Business Post* 48 (10) 411-420.
- 7) Chandrasekaran, V.K. 2013 *Flavour of Punjab*. *The Hindu* (Published on 6 March 2013).
- 8) Duke, J.A. and Wain, K.K. 1981. *Medicinal plants of the world*. Computer index with more than 85,000 entries, 3 vols.
- 9) Guy, C.L. 1990. Cold acclimation and freezing stress tolerance role of protein Metabolism. *Annu. Rev. plant physiol. Plant Mol. Biol.* 41:187-223.
- 10) Hayat, S.; Masood, A.; Yusuf, M.; Quazi, F. and Aquil A. 2009. Growth of Indian mustard in response to salicylic and under high temperature stress. *Brazilian J. of plant phy.* 21 (3).
- 11) Khan, M.A and Ungar, I.A. 1999. Effect of Salinity on the seed germination of *Triglochin maritima* under various temperature regimes. *Great Basin Naturalist* 59 (144-150)
- 12) Perry, L.M. 1980, *Medicinal plants of east and south east Asia* MTT Press, Cambridge.
- 13) Singh, J.; Laroche, A 1988. Freezing tolerance in plants; a biochemical overview. *Biochem Cell Biol* 66:650-657.
- 14) Zabator, F.J. 1962. Germination value: an index combining speed and completeness of Pine seed germination. *Forest science* 8:386-396.

Publish Research Article International Level Multidisciplinary Research Journal For All Subjects

Dear Sir/Mam,

We invite unpublished research paper.Summary of Research Project,Theses,Books and Books Review of publication,you will be pleased to know that our journals are

Associated and Indexed,India

- ★ International Scientific Journal Consortium Scientific
- ★ OPEN J-GATE

Associated and Indexed,USA

- DOAJ
- EBSCO
- Crossref DOI
- Index Copernicus
- Publication Index
- Academic Journal Database
- Contemporary Research Index
- Academic Paper Databse
- Digital Journals Database
- Current Index to Scholarly Journals
- Elite Scientific Journal Archive
- Directory Of Academic Resources
- Scholar Journal Index
- Recent Science Index
- Scientific Resources Database

Review Of Research Journal
258/34 Raviwar Peth Solapur-413005,Maharashtra
Contact-9595359435
E-Mail-ayisrj@yahoo.in/ayisrj2011@gmail.com
Website : www.isrj.net