



EMERGING TRENDS IN DNA MARKERS AND THEIR APPLICATIONS IN AQUATIC BIODIVERSITY

Bojjapalli Nagamani

Dept of Zoology, Osmania University Hyderabad.



ABSTRACT:

Biodiversity is essential for maintaining ecosystem services and each species performs a particular function in an ecosystem. It constitutes a unique global heritage and its conservation and utilization is of immediate concern. The increasing loss of aquatic biodiversity globally has led the researchers to intensify their efforts to produce a census of all aquatic biodiversity and to modernize the taxonomy. It is not feasible to catalogue the vast diversity by traditional methods based on morphological description so the researchers adopt analytical molecular technologies as an alternative to fill the gaps in phylogeny. In this direction, DNA markers have revolutionized the analytical power necessary to explore the genetic diversity. Among them mitochondrial DNA markers are being used extensively in aquatic biodiversity and conservation studies particularly in tracking invasive species, detection of cryptic species and identification of various life history stages. Focus has been given on mtDNA based markers such as DNA Barcoding, mini-DNA Barcoding and environmental DNA (eDNA) based meta-DNA Barcoding technologies useful for aquatic biodiversity studies.

KEY WORDS: aquatic biodiversity, cytochrome oxidase I (COI), DNA barcoding, environmental DNA (eDNA), molecular markers.

INTRODUCTION

Aquatic organisms have been the integral part of aquatic biodiversity and human beings have long been dependent on these resources for their food, health, environmental security and other commercial use. Biodiversity constitutes a unique global heritage and its protection and conservation is of immediate concern. Each species performs a particular function in an ecosystem. Aquatic biodiversity in marine and freshwater ecosystems is enormous and the quantity of explored diversity is far less than the actual diversity. The ancient ancestry extending into the past for more than 500 million years has allowed a vast span of time for evolutionary divergence and for the origin and extinction of major phyletic lines. Moreover, freshwater biodiversity has been experiencing alarming decline due to over exploitation of biota, habitat loss, anthropogenic pollution and other factors. As a result, several organisms were listed as endangered or threatened. The phylogenetic lineages based on the evolutionary relationships are still highly debatable inspite of repeated and continuing investigations using classical taxonomy. This scenario forced the researchers to adopt analytical molecular technologies as an alternative to fill the gaps in phylogeny and exploring the aquatic biodiversity. In this direction, DNA markers, genome mapping, microarrays and sequencing are proved to be the most relevant technologies. Molecular markers are potential tools to identify the genetic distinctiveness of individuals, populations or species. These markers have revolutionized the analytical strength necessary to investigate the genetic diversity. The development of DNAbased genetic markers created a huge progress in genetic studies. With these DNA markers, it is possible to observe and exploit genetic variation in the entire genome. Popular DNA markers in the aquatic biodiversity studies include:

Allozymes, mtDNA (Mitochondrial DNA) markers, RFLP (Restriction Fragment Length Polymorphism), RAPD (Random Amplified Polymorphic DNA), AFLP (Amplified Fragment Length Polymorphism), Microsatellites, SNP (Single Nucleotide Polymorphism) and EST (Expressed Sequence Tags) markers. Selection of appropriate genetic markers to assess the genetic diversity is the primary concern for any application in the field of aquatic biodiversity. Of late, mtDNA markers, especially DNA Barcoding and environmental DNA (eDNA) technologies are being gained greater attention from scientists with broad applications. Different types of markers used for aquatic biodiversity studies are depicted in Fig.1. In the process of evolution, every organism undergoes some genetic change at different magnitudes in reproductively isolated populations. Those changes enhance the capability of organisms to adapt to changing environment and are necessary for survival of the species. At the DNA level, genetic variation is mainly due to: base substitutions (Single nucleotide polymorphisms, SNPs); insertions or deletions of nucleotide sequences (Indels) within a locus; and the rearrangement of DNA segments around a specific locus. The present article collates information on molecular DNA markers with reference to their principles, potential power, applications and disadvantages in aquatic biodiversity studies. Emphasis has been given on recent technological advances in mtDNA based markers such as DNA Barcoding, mini-DNA Barcoding and environmental DNA (eDNA) based meta-DNA Barcoding technology.

MOLECULAR MARKERS

Allozyme Markers

Allozymes were among the earliest markers used in 1980's. These are the variant forms of an enzyme that are coded by different alleles at the same locus. They proved to be efficient both individually as well as in combination with other markers (RFLP, microsatellites or mitochondrial markers) for genetic stock assessment and revealing the genetic bottlenecks in various geographical regions. They are well known for estimating genetic differentiation and reproductive isolation, analysis of mating patterns coupled with Hardy-Weinberg pattern, which helps in determining the origin of an individual from a sample whether from a vast, randomly mating population with equilibrium genotype frequencies or from group of genetically distinct units. Allozymes revolutionized the studies on spawning aggregations by showing the homogeneity among heterogenous spawn both morphologically and life stage-wise over large geographic distances. Allozymes are not efficient with heterozygote deficiencies due to null (enzymatically inactive) alleles.

Microsatellite Markers

Microsatellites or Variable Number of Tandem Repeats (VNTR) are sequence repeats that range in size from 1 to 6 base pairs. Microsatellite loci typically exhibit elevated levels of length mutation facilitating high levels of heterozygosity in fish (ranging from 24-90%) which helps in stock identification. By using this technique, it is possible to compare the genetic diversity and population structure between wild and cultured stocks, which is useful for the researchers to improve species genetically by selective breeding and to design appropriate management guidelines for the conservation of a particular species. They provide valuable inferences while analyzing the genetic inheritance in the inter hybrid population and parentage analysis. Survey on genomic sequencing of channel catfish showed that microsatellites were found to represent 2.58% of the catfish genome. In most fish species, dinucleotide (AC)_n repeats are the most abundant forms of microsatellites. The disadvantage lies in the identification of microsatellite locus for its sequencing and it needs lot of investment and effort for primer designing. The mutation rates and patterns are also difficult to understand.

RAPD (Random Amplified Polymorphic DNA)

RAPD markers gain advantage with their arbitrary primers in several population study applications. RAPD can detect high levels of DNA polymorphism. The technique detects coding as well as non-coding DNA sequences, and most of the informative polymorphic sequences derived from repetitive (non-coding) DNA

sequences in the genome. RAPD markers have been widely used for species and strain identification in fishes and mollusks, genetic diversity and rate of gene flow in fish and analysis of genetic impact by environmental stressors. The major weakness of RAPD is its dominant mode of inheritance. Presence of paralogous PCR product (different DNA regions which have the same lengths and thus appear to be a single locus) limited the application of this marker.

RFLP (Restriction Fragment Length Polymorphism)

The RFLP technique is mainly based on DNA fragment length differences after digested genomic DNA with one or more restriction enzyme. Earlier these fragments were analyzed using Southern blot analysis by specific probes. Later, with the increasing number of “universal primers” developed from the flanking regions of fragments, it is replaced with PCR. RFLP is a co-dominant marker. It unable to detect point mutations, low polymorphic rates at most loci and can detect only large shifts in DNA fragment sizes. RFLP requires previous genetic information. Often this information is not available for many fish or other aquaculture species. Future use of RFLP will be to focus on analysis of Single Nucleotide Polymorphisms (SNP) residing within restriction sites.

AFLP (Amplified Fragment Length Polymorphism)

AFLP is a PCR-based, multi-locus fingerprinting technique that combines the strengths of RFLP and RAPD. Like RAPDs, it does not require any prior molecular information and thus applicable even to less studied fish species. By using AFLP, Co-dominant scoring is possible for well-characterized families. AFLP markers have been extensively used for genetic improvement programmes, genetic comparison of cultured and wild populations and other applications. The major weakness of AFLP marker is its dominant nature of inheritance. Also, it requires special equipment such as automated gene sequencers for electrophoretic analysis of fluorescent labels. Traditional electrophoretic methods can also be employed, but they require the use of radioactive labels or special staining techniques such as silver staining.

SNP (Single Nucleotide Polymorphism)

Single nucleotide polymorphism (SNP) describes polymorphisms caused by point mutations that give rise to different alleles containing alternative bases at a given nucleotide position within a locus. The PIC of SNP's is not as high as multi-allele microsatellites, but this limitation is balanced by their great abundance. The SNP's are inherited as co-dominant markers. Development of SNP-based genetic maps provides deep insights into genome-wide linkage study of sex determination and identification of loci on various chromosomes responsible for sex determination. SNP-based experiments showed direct evidence for polygenic sex determination in zebra fish. SNP's applicability to nonmodel organisms based on an annotated sequence of a model organism is a recently discovered application.

EST markers (Expressed Sequence Tags)

Expressed sequence tags (ESTs) are single-pass sequences generated from random sequencing of cDNA (complementary DNA) clones. It offers a rapid and valuable first look at the genes expressed in specific tissues under specific physiological conditions, or during specific developmental stages. ESTs are useful for the development of cDNA microarrays that allow analysis of differentially expressed genes to be determined in a systematic way, in addition to their great value in genome mapping. Recent development of radiation hybrid panels for widely cultivable species is a very good option for the application of EST markers in aquaculture.

Mitochondrial DNA markers (mtDNA)

These are strictly maternal inherited markers that contain faster mutation rate than the nuclear DNA. The recent application of mitochondrial markers is the identification of cryptic species and alien

species. mtDNA markers have emerged as one of the major tools for integrative taxonomy. mtDNA markers are useful for exploring the relationships among closely related species and for the identification of brood stocks. Different parts of the mitochondrial genome are known to evolve at different rates. Almost the entire mtDNA molecule is transcribed except for about 1-kb control region (D-loop) where replication and transcription of the molecule is initiated. In general, non-coding segments like D-loop exhibit elevated levels of variation relative to coding sequences such as cytochrome b (Cyt b) gene. Inventorying the “universal primers” elevated the use of these markers greatly in the field of aquatic biodiversity studies. Non-Mendelian inheritance and relatively minor proportion of mtDNA to total genome are the disadvantages for mtDNA markers. Gender biased inheritance (maternal) was also identified as another limitation to the validity of using mtDNA for genetic studies. Instances like back mutations (sites that have already undergone substitution are returned to their state), parallel substitutions (mutations occur at the same site in independent lineages), and rate of heterogeneity or mutational hot spots (large differences in the rate at which some sites undergo mutation when compared to other sites in the same region) also misleads the properties of mitochondrial markers.

DNA Barcoding

Before 2000s, various discrete units of mitochondrial genome were used in various applications without consistency among the scientific community. However, with the advent of DNA Barcoding technology using Cytochrome Oxidase I gene (COI/ Cox) by Noble laureate, PDN Hebert (2003), all the scientists come under one umbrella. This COI based technology is being used extensively in aquatic diversity studies for the identification of species, to examine phylogenetic relations, strains and hybrids as well as for species delimitation. This technology has solved many challenges in the field of classical taxonomy. It is proved as a successful tool for revealing cryptic species, calculating genetic divergence within and among the species, identification of Short Range Endemics (SRE's), etc. Moreover, it plays a crucial role in rapid documentation of α - taxonomy for several precious faunal groups before they lost and accelerate the rate of species discovery. However, the application of barcoding technology in taxonomy faced some controversies among scientists. Availability of universal primers facilitate the broad application of technology across all hierarchical levels of many groups of aquatic invertebrates and vertebrates including marine organisms. Food traceability is a recently emerged application of DNA Barcoding where it is more beneficial for quality assessment of aquatic products by controlling food fraudery activities, species substitutions and food authentication.

Mini DNA Barcoding

Mini DNA Barcoding is useful for degraded DNA samples and to identify the processed aquatic products. It focuses the analysis of shorter DNA fragments (e.g. 100–200 bp) within the full-length barcode. It is proved to be effective in obtaining DNA sequence information from specimens containing degraded DNA. Even short fragments of COI can be effectively used for species identification. With the advent of new sequencing technologies, this method gains much importance in determining the species composition of environmental samples, revealing prey-predatory relations, and identification of processed parts.

DNA meta Barcoding using environmental DNA (eDNA)

eDNA refers to the genetic material obtained from the environment which is in the form of whole microbial cells or shed off from multicellular organisms via metabolic waste, damaged tissue or sloughed skin cells. eDNA technology is based on the DNA released into environment in various forms like feces or excrements, fish slime, scrapped-off tissue cells and cells released after the death or decay of organism. It can be done even with low quantities of DNA. This technique is being widely used in determining the presence or absence of an organism in a particular aquatic environment which is critical to ecological management and conservation biology. It also estimates the abundance and biomass of a species in a

particular environment, creating the distributional maps, determining the feeding habits of endangered species and finding the prey-predatory relationships. eDNA monitoring has several advantages like increased sensitivity towards species detection, lower cost, extraction of DNA from multiple resources at a time and the possibility of targeting several taxa during amplification. Generation of sequencing data from environmental samples using eDNA is affordable. The major ecological applications of eDNA technology include tracking of invasive species and monitoring endangered species. eDNA technology helps in identifying the source species for whale meat, sturgeon eggs, shark fins and other high valued (and imperiled) species which are subjected to illegal trade. It is possible to track the presence of invasive species in an ecosystem before it gets established and cause harm to it. It can be applicable from the pond ecosystem level to the ocean ecosystem. Its application is expanded to many groups of animals including crustaceans mollusks, fishes and amphibians. Sometimes it requires past data for the analysis of eDNA. It is also possible to characterize diet contents of invasive species which helps in risk assessment of an ecosystem. Even small samples are sufficient for reliable identification. The main drawbacks of eDNA technology are: it is hard to obtain tissue samples of a single species from the whole sample; DNA fragments of interest often degrade faster in days to weeks beyond the level of detection in contemporary aquatic and marine ecosystems; false-positive or falsenegative detection rates; primer bias and variable eDNA concentrations; and expensive to adapt advanced PCR equipment (qPCR, ddPCR). Furthermore, different sets of marker combinations strengthen our ability in fishery management, rehabilitation and conservation of various species which are endemic and endangered due to destructive fishing practices and trade, over exploitation, habitat loss and restrictive distribution or even due to marine regression during glacial epoch. Allozyme and mitochondrial marker based studies revealed low levels of polymorphism, which pulls our discrimination power to assess stock structuring of freshwater fish. At this juncture microsatellites are proved to be the high polymorphic genetic marker than others. However, when dealing with highly divergent groups, microsatellites are less informative. Of late, various online and software analysis tools like e.g. MEGA V.7.0, Dna SP v6, POPGENE v1.32, Arlequin, MrBayes, XLSTAT, GenALEX etc. are available with fastest analytical power and giving reliable result for population data. Geneticists, conservationists and planners would take a strategic approach towards biodiversity monitoring by using molecular markers for sustainable management of aquatic biodiversity.

REFERENCES

1. Vitousek PM, Mooney HA, Lubchenco J, Melillo JM. Human domination of Earth's ecosystems. *Science*. 1997; 277(5325):494-499.
2. FAO. The State of World Fisheries and Aquaculture. FAO. 2004.
3. Pauly D, Christensen V, Guenette S, Pitcher TJ, Sumaila UR, Walters CJ et al. Towards sustainability in world fisheries. *Nature*, 2012; 418:689-695.
4. Srinivasan UT, Cheung WW, Watson R, Sumaila UR. Food security implications of global marine catch losses due to overfishing. *Journal of Bioeconomics*. 2010; 12(3):183-200.
5. Meffe GK. Conservation genetics and the management of endangered fishes. *Fisheries*. 1986; 11(1):14-23.
6. Liu ZJ, Cordes JF. DNA marker technologies and their applications in aquaculture genetics. *Aquaculture*. 2004; 238(1):1-37.
7. Hunter RL, Markert CL. Histochemical demonstration of enzymes separated by zone electrophoresis in starch gels. *Science*. 1957; 125(3261):1294-1295.
8. Singh RK, Lal KK, Mohindra V, Punia P, Sah RS, Mishra AK et al. Assessing genetic differentiation in geographic populations of *Labeo calbasu* using allozyme markers. *Biochem. Genet*, 2010; 48:760-778.
9. Richardson B, Baverstock PR, Adams M. Allozyme electrophoresis: a handbook for animal systematics and population studies. Academic Press, USA, 2012.
10. King DP, Ferguson A, Moffett IJ. Aspects of the population genetics of herring, *Clupea harengus*, around the British Isles and in the Baltic Sea. *Fisheries Research*. 1987; 6(1):35-52.

11. O'connell M, Wright JM. Microsatellite DNA in fishes. *Reviews in Fish biology and fisheries*. 1997; 7(3):331-63.
12. Lind CE, Evans BS, Knauer J, Taylor JJ, Jerry DR. Decreased genetic diversity and a reduced effective population size in cultured silver-lipped pearl oysters *Pinctada maxima*. *Aquaculture*. 2009; 286(1):12-9.
13. Christie MR, Marine ML, French RA, Waples RS, Blouin MS. Effective size of a wild salmonid population is greatly reduced by hatchery supplementation. *Heredity*. 2012; 109(4):254-60.
14. Kang JH, Kim YK, Park JY, An CM, Nam MM, Byun SG et al. Microsatellite analysis as a tool for discriminating an interfamily hybrid between olive flounder and starry flounder. *Genet. Mol. Res*. 2011; 10(4):2786-2794.
15. Pino-Querido A, Álvarez-Castro JM, Vera M, Pardo BG, Fuentes J, Martínez P et al. A molecular tool for parentage analysis in the Mediterranean mussel *Mytilus galloprovincialis*. *Aquaculture research*. 2015; 46(7):1721-1735.
16. Xu P, Wang S, Liu L, Peatman E, Somridhivej B, Thimmapuram J et al. Channel catfish BAC-end sequences for marker development and assessment of syntenic conservation with other fish species. *Animal genetics*. 2006; 37(4):321-326.
17. Bachtrog D, Agis M, Imhof M, Schlötterer C. Microsatellite variability differs between dinucleotide repeat motifs-evidence from *Drosophila melanogaster*. *Molecular Biology and Evolution*. 2000; 17(9):1277- 1285.
18. Welsh J, McClelland M. Fingerprinting genomes using PCR with arbitrary primers. *Nucleic acids research*. 1990; 18(24):7213-8.
19. Williams JG, Kubelik AR, Livak KJ, Rafalski JA, Tingey SV. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic acids research*. 1990; 18(22):6531-5.
20. Lerebours A, Cambier S, Hislop L, Adam-Guillermin C, Bourdineaud JP. Genotoxic effects of exposure to waterborne uranium, dietary methylmercury and hyperoxia in zebrafish assessed by the quantitative RAPD-PCR method. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*. 2013; 755(1):55-60.
21. Chu KH. Genetic variation in wild and cultured populations of the pearl oyster *Pinctada fucata* from southern China. *Aquaculture*. 2006; 258(1):220-227.
22. Praipue P, Klinbunga S, Jarayabhand P. Genetic diversity of wild and domesticated stocks of Thai abalone, *Haliotis asinina* Haliotidae, analyzed by single-strand conformational polymorphism of AFLP-derived markers. *Genetics and Molecular Research*. 2010; 9(2):1136-52.
23. Vignal A, Eggen A, Milan D, SanCristobal M. A review on SNP and other types of molecular markers and their use in animal genetics. *Genetics Selection Evolution*. 2002; 34(3):275-305.
24. Bradley KM, Breyer JP, Melville DB, Broman KW, Knapik EW, Smith JR et al. An SNP-based linkage map for zebrafish reveals sex determination loci. *G3: Genes, Genomes, Genetics*. 2011; 1(1):3-9.
25. Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE. Double digest RADseq: an inexpensive method for de novo SNP discovery and genotyping in model and non-model species. *PloS one*. 2012; 7(5):e37135.
26. Aoki JY, Kai W, Kawabata Y, Ozaki A, Yoshida K, Tsuzaki T et al. Construction of a radiation hybrid panel and the first yellowtail *Seriola quinqueradiata* radiation hybrid map using a nanofluidic dynamic array. *BMC genomics*. 2014; 15(1):165.
27. Brown WM, George M, Wilson AC. Rapid evolution of animal mitochondrial DNA. *Proceedings of the National Academy of Sciences*. 1979; 76(4):1967-1971.
28. Hebert PDN, Cywinska A, Ball SL. Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London B: Biological Sciences*. 2003; 270(1512):313-321.
29. Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PD. DNA barcoding Australia's fish species. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*. 2005; 360(1462):1847-57.
30. Hebert PDN, Gregory TR. The promise of DNA barcoding for taxonomy. *Systematic biology*. 2005; 54(5):852-859.

31. Kon T, Yoshino T, Mukai T, Nishida M. DNA sequences identify numerous cryptic species of the vertebrate: a lesson from the gobioid fish *Schindleria*. *Molecular Phylogenetics and Evolution*. 2007; 44(1):53-62.
32. Erpenbeck D, Weier T, De Voogd NJ, Wörheide G, Sutcliffe P, Todd JA et al. Insights into the evolution of freshwater sponges Porifera: Demospongiae: Spongillina: Barcoding and phylogenetic data from Lake Tanganyika endemics indicate multiple invasions and unsettle existing taxonomy. *Molecular Phylogenetics and Evolution*. 2011; 61(1):231-236.
33. DeSalle R, Egan MG, Siddall M. The unholy trinity: taxonomy, species delimitation and DNA barcoding. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 2005; 360(1462):1905-16.
34. Savolainen V, Cowan RS, Vogler AP, Roderick GK, Lane R. Towards writing the encyclopaedia of life: an introduction to DNA barcoding. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*. 2005; 360(1462):1805-1811.
35. Vrijenhoek R. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular marine biology and biotechnology*. 1994; 3(5):294-299.
36. Ivanova NV, Zemlak TS, Hanner RH, Hebert PD. Universal primer cocktails for fish DNA barcoding. *Molecular Ecology Resources*. 2007; 7(4):544-8.
37. Lobo J, Costa PM, Teixeira MA, Ferreira MS, Costa MH, Costa FO et al. Enhanced primers for amplification of DNA barcodes from a broad range of marine metazoans. *BMC ecology*. 2013; 13(1):34.
38. Tantillo G, Marchetti P, Mottola A, Terio V, Bottaro M, Bonerba E et al. Occurrence of mislabelling in prepared fishery products in Southern Italy. *Italian journal of food safety*. 2015; 4(3):152-156.
39. Khedkar GD, Tiknaik AD, Shinde RN, Kalyankar AD, Ron TB, Haymer D. High rates of substitution of the native catfish *Clarias batrachus* by *Clarias gariepinus* in India. *Mitochondrial DNA Part A*. 2016; 27(1):569-74.
40. Buhay JE. COI-like sequences are becoming problematic in molecular systematic and DNA barcoding studies. *Journal of Crustacean Biology*. 2009; 29(1):96-110.
41. Christie MR, Marine ML, Blouin MS. Who are the missing parents? Grandparentage analysis identifies multiple sources of gene flow into a wild population. *Molecular Ecology*. 2011; 20(6):1263-76.
42. Hajibabaei M, Smith M, Janzen DH, Rodriguez JJ, Whitfield JB, Hebert PD. A minimalist barcode can identify a specimen whose DNA is degraded. *Molecular Ecology Resources*. 2006; 6(4):959-964.
43. Shokralla S, Spall JL, Gibson JF, Hajibabaei M. Nextgeneration sequencing technologies for environmental DNA research. *Molecular ecology*. 2012; 21(8):1794-805.
44. Mitchell JK, Hellberg RS. Use of the mitochondrial control region as a potential DNA mini-barcoding target for the identification of canned tuna species. *Food Analytical Methods*. 2016; 9(10):2711-2720.
45. Ficetola GF, Miaud C, Pompanon F, Taberlet P. Species detection using environmental DNA from water samples. *Biology letters*. 2008; 4(4):423-425.
46. Rees HC, Gough KC, Middleditch DJ, Patmore JR, Maddison BC. Applications and limitations of measuring environmental DNA as indicators of the presence of aquatic animals. *Journal of Applied Ecology*. 2015; 52(4):827-831.
47. Takahara T, Minamoto T, Yamanaka H, Doi H, Kawabata ZI. Estimation of fish biomass using environmental DNA. *PLoS one*. 2012; 7(4):e35868.
48. Deiner K, Altermatt F. Transport distance of invertebrate environmental DNA in a natural river. *PLoS One*. 2014; 9(2):e88786.
49. Mächler E, Deiner K, Steinmann P, Altermatt F. Utility of environmental DNA for monitoring rare and indicator macroinvertebrate species. *Freshwater Science*. 2014; 33(4):1174-83.
50. Jerde CL, Mahon AR, Chadderton WL, Lodge DM. Sight-unseen detection of rare aquatic species using environmental DNA. *Conservation Letters*. 2011; 4(2):150-157.
51. Dejean T, Valentini A, Miquel C, Taberlet P, Bellemain E, Miaud C et al. Improved detection of an alien invasive species through environmental DNA barcoding: the example of the American bullfrog *Lithobates catesbeianus*. *Journal of applied ecology*. 2012; 49(4):953-959.

52. Takahara T, Minamoto T, Doi H. Using environmental DNA to estimate the distribution of an invasive fish species in ponds. *PLoS one*. 2013; 8(2):e56584.
53. Thomsen PF, Kielgast J, Iversen LL, Møller PR, Rasmussen M, Willerslev E et al. Detection of a diverse marine fish fauna using environmental DNA from seawater samples. *PLoS one*. 2012; 7(8):e41732.
54. Tréguier A, Paillisson JM, Dejean T, Valentini A, Schlaepfer MA, Roussel JM et al. Environmental DNA surveillance for invertebrate species: advantages and technical limitations to detect invasive crayfish *Procambarus clarkii* in freshwater ponds. *Journal of Applied Ecology*. 2014; 51(4):871-879.
55. Goldberg CS, Sepulveda A, Ray A, Baumgardt J, Waits LP. Environmental DNA as a new method for early detection of New Zealand mudsnails *Potamopyrgus antipodarum*. *Freshwater Science*, 2013; 32(3):792-800.
56. Takahara T, Minamoto T, Doi H. Effects of sample processing on the detection rate of environmental DNA from the Common Carp *Cyprinus carpio*. *Biological Conservation*. 2015; 183:64-69.
57. Darling JA, Mahon AR. From molecules to management: adopting DNA-based methods for monitoring biological invasions in aquatic environments. *Environmental research*. 2011; 111(7):978-988.
58. Muneer PA, Gopalakrishnan A, Musammilu KK, Basheer VS, Mohindra V, Lal KK et al. Comparative assessment of genetic variability in the populations of endemic and endangered Yellow Catfish, *Horabagrus brachysoma* Teleostei: Horabagridae, based on allozyme, RAPD, and microsatellite markers. *Biochemical genetics*. 2012; 50(3-4):192-212.
59. Molur S, Walker S. Conservation assessment and management plan for freshwater fishes of India. In Workshop Report by the Conservation Breeding Specialist Group, Zoo Outreach Organisation, Tamil Nadu, India, 1998.
60. Wang CF, Hsieh CH, Lee SC, Wang HY. Systematics and phylogeography of the Taiwanese endemic minnow *Candidia barbatus* Pisces: Cyprinidae based on DNA sequence, allozymic, and morphological analyses. *Zoological Journal of the Linnean Society*. 2011; 24;161(3):613-632.
61. Estoup A, Rousset F, Michalakis Y, Cornuet JM, Adriaamanga M, Guyomard R et al. Comparative analysis of microsatellite and allozyme markers: a case study investigating microgeographic differentiation in brown trout *Salmo trutta*. *Molecular Ecology*. 1998; 7(3):339- 353.
62. Shaw PW, Turan C, Wright JM, O'connell M, Carvalho GR. Microsatellite DNA analysis of population structure in Atlantic herring *Clupea harengus*, with direct comparison to allozyme and mtDNA RFLP analyses. *Heredity*. 1999; 83(4):490-499.
63. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular biology and evolution*. 2016; 33(7):1870-1874.
64. Librado P, Rozas J. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data.



Bojjapalli Nagamani
Dept of Zoology, Osmania University Hyderabad.