

Vol 5 Issue 11 August 2016

ISSN No : 2249-894X

---

*Monthly Multidisciplinary  
Research Journal*

*Review Of  
Research Journal*

Chief Editors

---

**Ashok Yakkaldevi**  
A R Burla College, India

**Ecaterina Patrascu**  
Spiru Haret University, Bucharest

**Kamani Perera**  
Regional Centre For Strategic Studies,  
Sri Lanka

Review Of Research Journal is a multidisciplinary research journal, published monthly in English, Hindi & Marathi Language. All research papers submitted to the journal will be double - blind peer reviewed referred by members of the editorial Board readers will include investigator in universities, research institutes government and industry with research interest in the general subjects.

**Regional Editor**

Dr. T. Manichander

Sanjeev Kumar Mishra

**Advisory Board**

Kamani Perera Regional Centre For Strategic Studies, Sri Lanka	Delia Serbescu Spiru Haret University, Bucharest, Romania	Mabel Miao Center for China and Globalization, China
Ecaterina Patrascu Spiru Haret University, Bucharest	Xiaohua Yang University of San Francisco, San Francisco	Ruth Wolf University Walla, Israel
Fabricio Moraes de Almeida Federal University of Rondonia, Brazil	Karina Xavier Massachusetts Institute of Technology (MIT), USA	Jie Hao University of Sydney, Australia
Anna Maria Constantinovici AL. I. Cuza University, Romania	May Hongmei Gao Kennesaw State University, USA	Pei-Shan Kao Andrea University of Essex, United Kingdom
Romona Mihaila Spiru Haret University, Romania	Marc Fetscherin Rollins College, USA	Loredana Bosca Spiru Haret University, Romania
	Liu Chen Beijing Foreign Studies University, China	Ilie Pinte Spiru Haret University, Romania
Mahdi Moharrampour Islamic Azad University buinzahra Branch, Qazvin, Iran	Nimita Khanna Director, Isara Institute of Management, New Delhi	Govind P. Shinde Bharati Vidyapeeth School of Distance Education Center, Navi Mumbai
Titus Pop PhD, Partium Christian University, Oradea, Romania	Salve R. N. Department of Sociology, Shivaji University, Kolhapur	Sonal Singh Vikram University, Ujjain
J. K. VIJAYAKUMAR King Abdullah University of Science & Technology, Saudi Arabia.	P. Malyadri Government Degree College, Tandur, A.P.	Jayashree Patil-Dake MBA Department of Badruka College Commerce and Arts Post Graduate Centre (BCCAPGC), Kachiguda, Hyderabad
George - Calin SERITAN Postdoctoral Researcher Faculty of Philosophy and Socio-Political Sciences Al. I. Cuza University, Iasi	S. D. Sindkhedkar PSGVP Mandal's Arts, Science and Commerce College, Shahada [ M.S. ]	Maj. Dr. S. Bakhtiar Choudhary Director, Hyderabad AP India.
REZA KAFIPOUR Shiraz University of Medical Sciences Shiraz, Iran	Anurag Misra DBS College, Kanpur	AR. SARAVANAKUMARALAGAPPA UNIVERSITY, KARAIKUDI, TN
Rajendra Shendge Director, B.C.U.D. Solapur University, Solapur	C. D. Balaji Panimalar Engineering College, Chennai	V.MAHALAKSHMI Dean, Panimalar Engineering College
Awadhesh Kumar Shirotriya	Bhavana vivek patole PhD, Elphinstone college mumbai-32	S.KANNAN Ph.D , Annamalai University
	Awadhesh Kumar Shirotriya Secretary, Play India Play (Trust), Meerut (U.P.)	Kanwar Dinesh Singh Dept.English, Government Postgraduate College , solan

More.....



## EFFECT OF WATER HARDNESS ON THE SURVIVAL RATE OF FRESH WATER FISHES.

**Shashikala Laxman Bhalkare**

Assit. Professor, Department of Fishery Science,  
Yeshawantrao Chavan College, Tuljapur, (M.S. India.)

### ABSTRACT

The aim of the present study was to investigate the effect of water source on survival rate %, growth performance, feed utilization, fish yield, economic evaluation and production of Nile tilapia (*Oreochromis niloticus*) monosex reared in earthen ponds. Nine earthen ponds were used and divided into three categories of three earthen ponds each. The average size of each pond was approximately 5200m<sup>2</sup>, 6000 monosex all male Nile tilapia were used in each pond and were stocked for 192 days. The fingerlings average weight was 4.38±0.03g/ fish, the fish were fed using a floating feed 25% crude protein, and were fed at a daily rate of 3% of their body weight. Results showed that body weight was increased significantly (P).

**KEYWORDS :** water Hardness , Fresh Water fishes , growth performance.

### INTRODUCTION

Nile tilapia, *Oreochromis niloticus* (Linnaeus) is one of the most important freshwater fish in world aquaculture. It is widely cultured in many tropical and subtropical countries of the world. Rapid growth rates, high tolerance to adverse environmental conditions, efficient feed conversion, ease of spawning, resistance to disease and good consumer acceptance make it a suitable fish for culture. Farmed tilapia production increased semi dramatically in recent years, increasing from 383,654mt in 1990 to 2,326,413mt in 2006. Tilapia has established a secure position in a number of water impoundments of India. But, its performance in open water ponds of the country has been discouraging over the years. For tilapia aquaculture is excessive reproduction and the resulting small size of the fish produced.

Egypt has suitable natural conditions for desert



aquaculture. Egypt has vast resources of groundwater. Fresh groundwater resources in Egypt contribute 20% to the potential water resources in Egypt. One of the groundwater resources is the Nile Valley and Delta system with the storage capacities of 200 billionm<sup>3</sup> and 300 billionm<sup>3</sup>, respectively. Oasis water in the west desert, Bahariya, Farafra, Dakhla, Kharga, and Siwa, were established from underground natural wells and springs.

With the prohibition of the establishment of fish farms on agricultural land, with the prohibition of the use of Nile water for fish farming, with increased competition for spaces adjacent to the lakes and sources of agricultural drainage water, despite its disadvantages, has caused the possession of new fish farm in the Nile. Valley of the most difficult things and out of reach. Hence the search for an

alternative to invade the desert, especially with the development of methods of fish farming and providing the requirements of education and with the provision of underground water of the highest purity with different salinity (fresh & brackish & marine) and where the trained professionals are available. In the hope to produce clean fish with improved quality and cheaper than other animal proteins we conducted the present research in a private fish farm located in the desert belonging to Noubaria Agricultural Development company (Ragab Farms) aiming to study the effect of water source on survival rate (%) growth performances, feed conversion ratio, protein efficiency ratio, annual fish yield and profitability Nile tilapia (*Oreochromis niloticus*) monosex commercial farming.

## MATERIAL AND METHODS

### Water source

Three types of water sources: fresh water, agricultural drainage water, and well water were compared in the present experiment. Water supplies were replaced three times during the experimental period (192 days).

### Experimental design

Nine earthen ponds (5200m<sup>2</sup>) were used in these experiment were divided into three categories of earthen ponds even three ponds represent one treatment (fresh water, drainage water and well water).

### Stocking density

6000 monosex all male Nile tilapia (*Oreochromis niloticus*) fingerlings of average weight (4.38±0.03g/fish) were stocked in each pond on April 11, 2007 and observed through October 19, 2007. The area of each pond 5200m<sup>2</sup>.

### Experimental fish

Fingerlings of all male Nile tilapia (*Oreochromis niloticus*) monosex were collected from Noubaria Agricultural Development Company (Ragab Fish Hatchery) and were over wintered in earthen ponds to provide suitable fingerlings for the beginning of the growing season. All ponds in these experiment was sampled monthly using a cast net method. Sample sizes were 1% of the stocked numbers and the average individual fish weight was calculated to determine growth rates. Then, with these calculations, the feed amounts were adjusted for the following month.

### Experimental diet

The floating commercial diet used in this experiment was fed at a daily rate of 3% of the fish body weight by using self feeders. The ingredients of the commercial diet used in the experiment is presented in Table 1. The dietary composition of vitamin and mineral premix is listed in. Fish were fed a floating ration for 6 days per week. Feeding rate was adjusted monthly based upon the calculated biomass of fish obtained through the monthly sampling and assumption of 100% survival.

**Table 1: Composition and proximate composition of control and the experimental diets for feeding Nile tilapia.**

Ingredients (%)	Diet
Fish meal	10
Meat meal	10
Soybean meal	16
Wheat bran	20
Yellow corn meal	38
Corn oil	3
Bone meal	1
Vitamin mixture	1
Minerals mixture	1
Proximate composition (%)	
Dry matter	90.65
Crude protein	25.12
Ether extract%	5.9
Ash	7.41
Crude fiber	4.4
NFE4	56.17
Goss energykcal/g	4.3

### Water quality

**a. Physical parameters:** Water temperature °C was determined at every days in the experiment.

**b. Chemical parameters:** Samples for determination of Dissolved Oxygen (DO) were immediately fixed after sampling and DO concentration was determined according to Winkler's technique. Methods described by Golterman were used in determination of ammonia. Also pH was measured by digital pH meter (Orion model 720 A, s/No 13602) in all experiments.

### Chemical analysis of the commercial of diet

Chemical analysis of the commercial diet used in the experiment was done according to [10] as shown in Table 1.

### Growth parameters and statistical analysis

Data on growth, feed utilization, survival rate and proximate and chemical composition of whole fish body were subjected to one-way ANOVA. To locate significant differences between fish size within different water resources of pond. Duncan's multiple rang test was done. All percentages and ratio were transformed to arcsine values prior to analysis.

## RESULTS AND DISCUSSION

### Experimental diet

The commercial diet used in the present experiment contained 25% CP and 4.3kcal/g gross

energy (Table 1). Although there are large variations in the data available about the optimum protein level for tilapias which range between 20 and 40% crude protein practical diets as low as 25% protein was successfully used for rearing monosex tilapia.

**Table 2: Water quality parameters (Mean±SE) of fresh water, drainage water well water during 8 months of the experimental period (April to November 2008).**

*Duration Months	April	May	June	July	August	September	October	November
<b>Fresh Water</b>								
Ammonia mg/l	0.09±0.01	0.11±0.01	0.12±0.01	0.12±0.01	0.12±0.01	0.11±0.01	0.09±0.01	0.12±0.01
Ph	7.98±0.13	7.98±0.13	8.00±0.13	8.00±0.13	8.00±0.13	8.00±0.13	8.00±0.13	7.98±0.13
D Oxygen mg/l	6.58±0.92	6.83±0.61	6.82±0.56	7.33±0.09	7.08±0.53	7.33±0.09	7.37±0.09	8.33±0.09
Temperature	25.4±0.12	26.2±0.14	28.9±0.67	29.6±0.43	30.3±0.45	28.4±0.62	26.9±0.13	24.13±0.53
<b>Drainage Water</b>								
Ammonia mg/l	0.10±0.01	0.10±0.01	0.11±0.01	0.11±0.01	0.13±0.01	0.13±0.01	0.12±0.01	0.11±0.01
Ph	8.01±0.13	8.03±0.13	8.04±0.13	8.06±0.13	8.10±0.13	8.09±0.13	8.06±0.13	8.06±0.13
D. Oxygen mg/l	6.56±0.90	6.80±0.60	6.82±0.58	7.31±0.08	7.07±0.52	7.33±0.09	7.34±0.08	8.33±0.09
Temperature °C	25.98±0.1	26.60±0.2	29.04±0.8	30.29±0.4	30.65±0.5	28.48±0.7	27.28±0.1	24.23±0.5
<b>Well Water</b>								
Ammonia mg/l	0.06±0.01	0.07±0.01	0.08±0.01	0.08±0.01	0.10±0.01	0.09±0.01	0.08±0.01	0.06±0.01
Ph	8.00±0.13	8.02±0.13	8.04±0.13	8.05±0.13	8.10±0.13	8.08±0.13	8.04±0.13	8.00±0.13
D Oxygen mg/l	6.07±0.52	6.22±0.58	6.31±0.08	6.52±0.60	6.56±0.90	6.34±0.08	6.33±0.09	6.33±0.09
Temperature °C	9.94±0.12	30.82±0.14	0.85±0.67	32.46±0.4	3.63±0.43	33.38±0.62	32.88±0.1	31.63±0.53

**Table 3: Effect of water source on growth performance of Nile tilapia monosex during the experimental period (days)**

Item	Body weight (g)		Daily gain (g)	SGR	FCR	PER
	Initial	Final				
Fresh water	4.38±0.03	354.2±14.42b	1.82±0.07b	2.27±0.02b	2.87±0.00a	1.38±0.00c
Drainage water	4.35±0.08	320.2±19.47b	1.64±0.10b	2.22±0.02b	2.80±0.00c	1.42±0.00a
Well water	4.36±0.07	472.3±2.92a	2.44±0.01a	2.43±0.00a	2.83±0.00b	1.40±0.00b
LSD(P<0.05)*	NS	42.5	0.22	0.06	0.02	0.01

Means in the same column having different letters are significantly different (P<0.05).

**Table 4: Effect of water source, pond sizes stocking density on yield and survival rate (%) of tilapia monosex**

Item	Total yield		Survival %
	Kg	%	
<b>Water source</b>			
a) Fresh water	2128±86.43 <sup>b</sup>		98.53±0.05 <sup>a</sup>
b) Drainage water	1921.8±116.96 <sup>b</sup>		95.05±0.3 <sup>b</sup>
c) Well water	2837.7±617 <sup>a</sup>		98.31±0.42 <sup>a</sup>
LSD (P<0.05)	255.01		0.91

Means in the same column having different letters are significantly different (P<0.05).


Overall means of water Dissolved Oxygen (DO) throughout the present experiment were

7.20±0.37mg DO/l for fresh water, 7.19±0.36mg DO/l for drainage water and 6.33±0.36 mg DO/l for well water. The fluctuation of water Dissolved Oxygen (DO) showed that the maximum values of DO were obtained in November for the fresh and drainage water and August in well water, however, the lowest values were in April. In general, dissolved oxygen levels were within the high standards and higher than cited by Boyd for good production of tilapia (4.20 to 5.90mg DO/l) in aquaculture ponds. One of the most important environmental factors is dissolved oxygen. It is considered a limiting factor for success or failure in intensive culture. An excellent aquaculture attribute of tilapia is their tolerance to low dissolved oxygen concentration. The dissolved oxygen content in earthen ponds depends on the pond water temperature, fish biomass and rate of water exchange. Chervinski reported that *O. niloticus* survived short term exposure to 0.1mg DO/ l. However, Collins observed in a review on oxygen concentration of various studies, that growth rate of non-salmonid fish was increasingly depressed as dissolved oxygen fall below 50% saturation. Rappaport reported that growth of carp was reduced by predawn dissolved oxygen less than 25% saturation. Tichert-C & Green compared the growth of tilapia monosex in earthen ponds aerated or unaerated at 10 or 30% saturation of dissolved oxygen. They found that tilapia production and final weight were significantly greater in aerated ponds than unaerated ponds.

The water pH values throughout the present experiments ranged between 8.00±0.13 and 8.10±0.13 with an overall mean of 8.04±0.13 in fresh water and ranged between 8.01±0.13 and 8.10±0.13 with an overall mean of 8.05±0.13 in drainage water and ranged between 7.98±0.13 and 8.01±0.13 with an overall mean of 8.00±0.13 in well water. The fluctuations of pH reach the highest value of 8.10±0.13 during August in fresh and drainage water and 8.01±0.13 in well water. The results showed that the present pH values are suitable. For rearing tilapia monosex in earthen ponds. Johnson recommended the range of pH 6.5 to 9.0 for most of freshwater fish species.

The water un-ionized ammonia (NH<sub>3</sub>) throughout the present experiments ranged between 0.09±0.01 and 0.12±0.01 with an overall mean of 0.11±0.01 in fresh water and ranged between 0.10±0.01 and 0.13±0.01 with an overall mean of 0.11±0.01 in drainage water and ranged between 0.06±0.01 and 0.10±0.01 with an overall mean of 0.077±0.01 in well water. The fluctuations of un-ionized ammonia reach the highest values of 0.13mg/l during August. Unionized ammonia concentrations in the experimental ponds generally remained below levels which would cause chronic toxicity problems in tilapia. Tilapia is more tolerant to elevated levels of ammonia than more other sensitive species such as salmonids. Some tilapias have been shown to acclimate to higher levels of ammonia after chronic exposure to low levels. Johnson showed that levels of un-ionized ammonia which may adversely affect growth in tilapia range from 1mg/l to 2mg/l ammonia where temperature and pH are within normal range.

### Growth performance of tilapia monosex



**Mean weight:** Results of the present study showed that the mean weights at all rearing intervals different significantly ( $P < 0.05$ ) during all the experimental periods (Figure 1). Averages of fish body weights for fresh water, drainage water and well water were found to be 23.16, 18.66 and 25.16g, respectively after the 1st month of stocking. The statistical evaluation of results indicated that live weights at this period increased significantly ( $P < 0.05$ ) with using well water. A similar trend was also observed in fish body weights during the other growing periods. At harvest average body weight of fish stocked at well water was significantly ( $P < 0.05$ ) higher than that of fish stocked in fresh or drainage water, which indicates that weights fish were decreased in fresh and drainage water with increasing for used well water at harvest were 354.17g, 320.17g and 472.33g for fresh, drainage and well water, respectively. This significant advancement in fish body weights with increasing at higher temperature of water advocated by Azaza reported that the final mean weight was significantly higher at 26 and 30 °C than at 22 and 34 °C. This finding agree with our results.

**Average daily gain (ADGg/day):** Results presented in Table revealed that water sources, affected significantly ( $P < 0.05$ ) ADG during all experimental periods tested (30, 60, 90, 120, 150, 180 and 210 days after start). In general these results indicated that the well water favored significantly ADG of the tilapia monosex in intensive culture system. The results of this point were in agreement with those found by who grew *O. niloticus* from 49g to 271g in 122 days (1.4%/day). Siddiqui found that ADG of tilapia *O. niloticus* reared for 98 days at different water exchange in outdoor concrete tanks was 1.06g/day at 30% dietary crude protein. In the present study the average daily gain was higher with 25% crude protein at all treatments. However, the optimal feeding rate depends on fish size and water temperature.

**Specific growth rate (SGR%):** Results presented in Table, revealed that water sources, affected significantly ( $P < 0.05$ ) SGR% during all experimental periods tested (30, 60, 90, 120, 150, 180 and 205 days after start). In general these results indicated that the well water, favored significantly ( $P < 0.05$ ) SGR% of the tilapia monosex in intensive culture system. During all tested experimental periods tested (30, 60, 90, 120, 150, 180 and 205 days after start) SGR% increased significantly ( $P < 0.05$ ) in almost linear manner in the well water than fresh and drainage water. In the present study SGR% values in case of well water continuously higher than fresh or drainage water in all experimental periods. This may be due to the higher temperature of the well water (average 31.94 °C) compared to 27.47 and 27.81 °C for fresh and drainage water, respectively. The results obtained in SGR% are in agreement with those found by Eid & El Denasoury who indicated that increasing temperature from 16 °C to 27 °C improve growth



rate of Nile tilapia, which using well water.

**Feed conversion ratio (FCR):** Results presented in table, show that there were significant ( $P < 0.05$ ) effects of water sources on FCR, feed conversion ratio was observed at harvest was 2.87 at fresh water, followed by 2.83 at well water and 2.80 at drainage water and 2.94 for 1700m<sup>2</sup> followed by 2.89 for 4000m<sup>2</sup> followed by 2.75 for 5200m<sup>2</sup> and was 2.57 for 6000 fish/acre, 2.75 for 8000 fish/acre and 2.78 for 10000 fish/ acre. The analyses of variance of the FCR values are presented in table . The FCR is affected by the physiological state of the fish, environmental condition. Lovshin found that FCR for all male tilapia in earthen ponds was higher (4.3) than when compared with all male and female tilapia in earthen ponds (FCR=7.2). While, fish growth is affected by the amount of feed consumed and the efficiency of assimilation.

**Protein Efficiency Ratio (PER):** Results of Protein Efficiency Ratio (PER) are presented in table, There were significant ( $P < 0.05$ ) effects of water sources, on PER, it improved significantly ( $P < 0.05$ ) with each increase in pond sizes and decrease stocking density throughout the experimental periods. The best PER observed at harvest was 1.42 with drainage water, followed by 1.40 at well water and 1.38 at fresh water confirmed that when protein supply is appropriate (400-500g protein/kg feed for percid fish), different lipid contents in feeds do not have an impact on the rearing results of pikeperch.

**Fish Survival Rate:** Results in table showed that survival rates were changed significantly ( $P < 0.05$ ) by water resources, in fresh and well water were insignificantly ( $P < 0.05$ ) different but survival of the fish in drainage water was 95.05% which was less than survival rates in both fresh water and well water indicating the probable effect of some faction of water quality.

**Fish Yield:** Results of (Table 4) show fish yield (kg) per acre as affected by water sources. Results revealed that total yield increased significantly ( $P < 0.05$ ) with well water. The total production was found to be 133.34% and 90.31 % for well water and drainage water, respectively, while it was found to be 76.33% and 68.84% for 4000 m<sup>2</sup> and 5200m<sup>2</sup>. The results of the present experiment were similar to those of who showed that the net yield of tilapia monosex in earthen ponds was 16750Kg/ha (7035.0kg/acre) after 100 days of stocking of 80.000 fish/ha, (33600 fish/acre, 8 fish/m<sup>2</sup>) on the other hand indicated that increasing temperature from 16 °C to 27 °C improved growth rate of Nile tilapia. Watanabe found that growth rates generally increase with increasing temperature and were markedly lower at 22 °C and well water is the best because the temperature constant through the year and the best quality of the water. Zhang found the higher yield obtained in small pond sizes because the bigger ponds with greater surface area were more difficult to manage and often resulted in lower fish yields. All fish species are characterized by an ideal range of temperature in which they show their maximum growth. Several studies have been reported that the specific water temperature range showed the faster growth in Pikeperch, Sander lucioperca at 20 °C to 25 °C. Low temperature causes sluggishness by retarding the digestion speeding of fish. Some researchers have found that the digestion rate has been increased as the temperature increases. Environmental temperature is one of the most important ecological factor which also influence the behavior and physiological process of aquatic animals [45]. One of the major advantages of groundwater sources is their constant temperature throughout the year. Shallow sources of groundwater approximate the mean air temperature of the area. The chemistry of groundwater is directly dependent on the geology of the area surrounding the source. In limestone areas, groundwater is hard, and high in calcium and carbon dioxide. In areas of granite formation, the groundwater tends to be soft, low in dissolved minerals and carbon dioxide. As will be discussed later, there are advantages and disadvantages to both, emphasizing the need for early extensive water quality testing. Water temperature has substantial effect on fish metabolism. In response to decreasing of water temperature the enzyme activity of

tissues have been increased. Velmurugan have investigated that histopathological and tissue enzyme changes of *C. gariepinus* exposed to nitrite when water temperatures changes from 27 °C to 35 °C. In a stressful and unfavorable environmental condition GPT and GOT may increase in blood serum . In the present study serum GPT and GOT level were affected by different water temperature. Serum GPT and GOT amount in different fish fed at 20 °C are comparatively lower than those of fish fed at 16 °C and 24 °C experiments (Tables 1-3). These results indicated that 20 °C may be a favorable water temperature for better growth of 16g juvenile Korean rockfish.

#### REFERENCES:

1. AS Fineman-K, AS Camacho (1987) The effects of supplemental feeds containing different protein: energy ratios on the growth and survival of *Oreochromis niloticus* (L.) in brackish water ponds. *Aquaculture Research* 18(2): 139-149.
2. Siddiqui AQ, MS Howlader AA, Adam (1988) Effects of dietary protein levels on growth, feed conversion and protein utilization in fry and young Nile tilapia, *Oreochromis niloticus*. *Aquaculture magazine* 70(1): 63-73.
3. El-Saidy DMS, Gaber MM (2005) Effect of dietary protein source and feeding frequency on growth performance, production traits and body composition of Nile tilapia, *Oreochromis niloticus* (L) cultured in concrete tanks. *Egyptian J Nutrition and feeds* 8(1): 1077-1089.
4. Collins G (1984) Fish growth and lethality versus dissolved oxygen. *Proceeding of the 1984 Specialty Conference on Environmental Engineering, 25-27 June 1984, ASCEJ, Los Angeles, CA, USA, pp. 750- 755.*
5. Johnson SK (1986) Ground water: Its quality characteristics aquaculture. *Proc. Texas fish farming conf., Texas A&M University, USA.*
6. Brian JB, James SD (1994) Effects of feeding frequency and handling on growth and mortality of cultured walking catfish *Clarias fuscus*. *Journal of the World Aquaculture Society* 25(2): 175-182.
7. Andras R, Istvan C (2008) Effect of feeding regime and temperature on on-growing results of pikeperch (*Sander lucioperca* L.). *Aquac Res* 39(8): 820-827.
8. Turker Ali (2009) Effect of photoperiod on growth of trout (*Oncorhynchus mykiss*) in cold ambient sea water. *Israeli journal of aquaculture-Bamidgeh* 61(1): 57-62.

# Publish Research Article

## International Level Multidisciplinary Research Journal For All Subjects

Dear Sir/Mam,

We invite unpublished Research Paper, Summary of Research Project, Theses, Books and Books Review for publication, you will be pleased to know that our journals are

### Associated and Indexed, India

- ★ Directory Of Research Journal Indexing
- ★ International Scientific Journal Consortium Scientific
- ★ OPEN J-GATE

### Associated and Indexed, USA

- DOAJ
- EBSCO
- Crossref DOI
- Index Copernicus
- Publication Index
- Academic Journal Database
- Contemporary Research Index
- Academic Paper Database
- Digital Journals Database
- Current Index to Scholarly Journals
- Elite Scientific Journal Archive
- Directory Of Academic Resources
- Scholar Journal Index
- Recent Science Index
- Scientific Resources Database

Review Of Research Journal  
258/34 Raviwar Peth Solapur-  
413005, Maharashtra  
Contact-9595359435

E-Mail-ayisrj@yahoo.in/ayisrj2011@gmail.com