



## ARBUSCULAR MYCORRHIZAL COLONIZATION IN *FICUS BENGHALENSIS* OF FAMILY MORACEAE


**Dr. Savita B. Wankhede<sup>1</sup> & Dr. R. M. Mulani<sup>2</sup>**

<sup>1</sup>Assistant Professor, Rajiv Gandhi Mahavidyalaya Mudkhed,  
Dist.Nanded, Maharashtra, India.

<sup>2</sup>UGC-SAP and DST-FIST sponsored school of life sciences,  
Swami Ramanand Teerth Marathwada University Nanded,  
Maharashtra, India.

### ABSTRACT

Arbuscular mycorrhizal fungi play an important role in the mobilization nutrients and enhancing plant growth. It maintains the intimate link between the plant roots and soil. Present work deals with Arbuscular Mycorrhizal colonization in *Ficus benghalensis* and isolation of resting spores from rhizospheric soil of *Ficus benghalensis* from fields of swami Ramanand Teerth Marathwada University Nanded and were analyzed by using wet sieving and decanting method suggested by Gerdman and Nicolson method (1963). The spores were analyzed *Glomus* sp. and *Aculospora* sp. Whole mount of root was analyzed for the root colonization by using the method suggested by Phillips and Hymen (1970). The % colonization was 80 to 90% and the root colonization showed rounded, elongated Vesicles and Arbuscles.



**KEYWORDS:** Arbuscular Mycorrhizal fungi, Root colonization

### INTRODUCTION

*Ficus benghalensis* is used for treatment of fever, ulcers, inflammation vomiting and leprosy. The latex is useful in piles, nose diseases and gonorrhoea. The root of *Ficus benghalensis* is useful in dysentery, inflammation of liver. Milky juice is used for rheumatism, pains and Seeds are cooling and tonic in nature. The bark is used for diabetes and leucorrhoea also used as astringent and tonic. This tree found in everywhere.

**Classification: (Tripathi and Kumar, 2015).**

Kingdom: Plantae

Division: Magnoliophyta.

Class: Magnoliopsida

Order: Utricales

Family: Moraceae

Genus: *Ficus*

Species: *benghalensis*

German Botanist Frank (1885) coined the term mycorrhizae for the first time to designate the symbiotic relationship between the fungi and plant roots. Since then scientists started exploiting them for the welfare of mankind. The term 'mycorrhiza' in its broadest sense is the non-pathogenic association of fungi and the roots of higher plants. The root- fungus association is symbiotic and the whole association is being considered as a "functionally distinct organ" involved in mineral nutrient uptake from the soil. (Kar, 1993). Mycorrhizal fungi are having intimate association with roots of higher plants forming a symbiotic relationship providing nutrients to the plants. The Arbuscular Mycorrhizal diversity in herbaceous vegetation medicinal plants, in halophytes plants have been investigated by many workers [Bagyaraj, D. J. (2014) Kannan, K. and Lakshminarashiman, C. (1988) Kumar., et. al (2013). Mulla, R. M et. al., (1994) Mulani., R. M et. al., (2004) Mulani, R. M and Waghmare, S. S. (2012). Mulani, R. M and Prabhu, R. R. (2002). Parameswaran, P and Augustine, B.(1988). Isolation and identification of arbuscular mycorrhizal fungi from agricultural fields of Vietnam investigated by (Sasvari et.al., 2012). Growth and biomass of *Piper longum* L was increased with inoculation of arbuscular mycorrhizal fungi. (Seema and Rajkumar,2015). Essential oil production, nutrient uptake and root colonization in basil was increased with inoculation arbuscular mycorrhizal fungi. (Mirhassan et.al.,2010).

## MATERIALS AND METHODS.

### Isolation of spores by using wet-sieving method. (Gerdman and Nicolson; 1963).

Spore extraction is involved in three sub steps such as wet-sieving, sedimentation, flotation. Mix 5 gm of soil in 250 ml of lukewarm water in a beaker until all aggregates disperse to a uniform suspension. Allow the heavier particles to settle down. Filter the suspension through 710 µm sieve to remove large organic matter and roots. Then solution was sieved through series of sieves i.e. 710 µm, 210 µm, 150 µm, 75 µm, 45 µm and 25 µm respectively. Content of each sieve i.e. 210 µm, 150 µm, 75 µm, 45 µm and 25 µm was taken separately on blotting paper in petriplate and This petriplate was observed under stereo zoom binocular microscope.

### Percentage of root colonization. (Phillips and Hayman, 1970).

Young root segments were taken in test tube adding 10% KOH and it autoclaved at 15 lbs for 1 hr. After 10 minute 10% KOH was removed from test tube then root segments were washed under tap water with 2 to 3 times. Then 10 ml 1N HCL was added and were kept for 5 minute for neutralization of root tissue. Then HCL was removed and washed the root segments 2 to 3 times with tap water. After 30 minute root segments stained with cotton blue and kept for 24 hrs. After 24 hrs root segments mounted on slide with Acetic acid - glycerol (1:1v/v). Seal the corners of the cover slip with DPX, root colonization was observed under compound microscope. Then % of Arbuscular Mycorrhizal fungal colonization calculated by using this formula


$$\text{Percent of mycorrhizal colonization} = \frac{\text{Number of root segments colonized}}{\text{Total number of root segments examined}} \times 100$$

## RESULTS AND DISCUSSION:

The roots of *Ficus benghalensis* showed 80 to 90% Mycorrhizal colonization and the rounded, vesicles were prominent. The rhizospheric soil was screened for spore density and population. The spore density were recorded as 265 spores per 100gm of soil and The spore population mainly consist of different species of Arbuscular mycorrhizal such as mainly dominated by *Glomus* sp. The other species were *Aculospora* and *Gigaspora*. Similar observation made by Sasvari et. al., (2012) in their studies highest number of spores found in the tomato and peanuts at agricultural field of Vietnam. The roots of *Aloe vera* showed 90 % root colonization and spore density was recorded as 250 spores per 100 gm of soil. Such observation were made by Mulani and Waghmare,(2012). The presence of large number of spore with varied population of spores indicated their universal occurrence in the soil of

university campus. Such observations were made by Mulani and Prabhu. (2002), Mulani et.al., (2004), Prabhu(2002) and Sathe (2005).Mulani and Prabhu had observed highest count of chlamydospores occurring in the root zone soil of *Dipcadi saxorum*. The murmy soil with moisture % and low humidity with high temperature fevers more chlymadospre formation. Similar observations were made by Harinikumar and Bagyaraj (1988) and Bagyaraj (1995) in tropical soil. Recently Pawar and Kakde (2012) have carried out the studies on the AMF associated with some medicinal plants from Mumbai region. They reported eight different species of *Glomus* namely *G. aggregatum*, *G. Boreale*, *G. fasciculatum*, *G. geosporum*, *G. heterosporum*, *G. segmentatum*, *G. tortuosum*, *G.radiatum* associated with the selected medicinal plants.

Root colonization of *Ficus benghalensis* showing in fig: a, b, c, d, e, f, Magnified view of oval and rounded vesicles, Hyphae seen in whole mount of root of *Ficus benghalensis* Spores were isolated from rhizospheric soil of *Ficus benghalensis* i.e *Glomus* sp.fig. a (100x). *Acaulospora* sp.(fig b.100x).





**REFERENCES:**

- Bagyaraj, D. J. (2014). Mycorrhizal fungi. Proc Indian Natn Sci Acad. 80(2): 415-428.
- Bagyaraj, D. J. (1995). Influence of agricultural practices on vesicular arbuscular mycorrhizal fungi in soil. Journal soil biol. Ecol. 15(2):109-116.
- Gerdman, J. W. and Nicolson, T. H. (1963). Spores of mycorrhizal Endogone species extracted from the soil by wet sieving and decanting. Trans. Br. Mycol. Soc. 46:235-244.
- Harinkumar, K.M and Bagyaraj, D.J. (1988). The effect of season on vesicular Arbuscular Mycorrhiza of Leucaena and Mango in a semiarid tropic. Arid soil Res. Rehabi.7:139-143.
- Kar, A.K. (1993) College Botany II New Central Book Agency, Calcutta. 507.
- Imam, M and Akter S(2011).Musa paradisiacal L. and Musa sapientum L. A Phytochemical and pharmacological review. Journal of Applied pharmaceutical science,01(05);pp.14-20.
- Kannan, K. and Lakshminarashiman, C. (1988). Survey of VAM of maritime strand plants of Po Calimere. In-First Asian conference on Mycorrhizae, C.A.S. in Botany, Madras. 29(31): 53-55.
- Kumar., A. Chhavi, M. and Aggrawal, A. (2013).Biodiversity of Endophytic mycorrhizal fungi associated with some medicinal plants of Himachal Pradesh. Asian J. of Adv. Basic sci. 1 (1): 26-29.
- MirHassan, R., Abbas. H., Mohsen B., Younes., R and Fatemeh., S.(2010) Effects of arbuscular mycorrhizal (AM) fungi on growth, essential oil production and nutrients uptake in basil. Journal of Medicinal Plants Research Vol.4(21), pp. 2222-2228.
- Mulla, R. M. and Kanade, A. M. (1994). VAM Mycorrhizal colonization in grasses of Bombay. J. Rayat Shikshan Sanstha Satara : 56-65.
- Mulani, R. M, Prabu, R. R. and Dinkaran, M. (2004).Occurance of vesicular Arbuscular Mycorrhizaa (VAM) in the roots of phylanthusfraternus Webster. Mycorrhiza News. 14 (2):11-14.
- Mulani, R. M and Waghmare, S. S. (2012). Assessment of occurrence of Thermo tolerant Arbuscular Mycorrhizal Fungi in the Roots and Rhizospheric spoil of Aloe vera (.).Burn.f. Online international journal interdisciplinay research journal. 2(4): 22-27.
- Mulani, R. M. and Prabhu, R. R. (2002). A seasonal variation in Arbuscular Mycorrhizal (VAM) colonization in the roots of Dipcadisaxorum Blatt and chlamyospores in the rhizosperic soil from Mumbai. J. sol. Biol. & ECOL. 20 (172): 47-50.
- Parameswaran, P and Augustine, B. (1988). Distribution and ecology of VAM in a scrub jungle. In-First Asia conference on Mycorrhizae, C. A. S. in Botany, Madras. 29(31): 91-99.
- Pawar, J. S and Kakde, U. B (2012).Study of arbuscular Mycorrhiza associated with some important medicinal palnts suburban area of Mumbai. Online international journal interdisciplinary research journal. II. 116-127.
- Phillips, J. M. and Hayman, D. S. (1970). Improved procedure for clearing roots and staining parasitic and vesicular arbuscular mycorrhizal fungi for rapid assessment of infection. Trans. Br. Mycol. Soc. 55: 152-160.
- Prabhu, R. R. (2002).Survey of soil of Mumbai and Adjoining areas for native VAM and their multiplication and effect of their inoculation on local crops as biofertilizers. A Ph.D thesis submitted to Mumbai University.
- Sasavari, Z, Magurno. F, Galanics, D, Hang T. T, Hong Ha. T. T, Luyen .N. D, Huong .L and Posta .K (2012). Isolation and identification of Arbuscular Mycorrhizal Fungi from agricultural fields of Vietnam .American Journal of plant sciences 3, 1796-1801.
- Sathe, V. D.(2005). Assessment of arbuscular mycorrhizal status in the soil on some forest plateaus of the Western ghats of Maharashtra. A Ph.D thesis submitted to the Mumbai University.
- Seema, H. S and Rajkumar H. G. (2015) Effect of arbuscular mycorrhizal fungi on growth and biomass enhancement in piper longum L.(Piperaceae). Int. J. Curr. Microbiol. App. Sci (2015) 4(1): 11-18.